

# Specimen Label



# Rodeo®

## Herbicide

For control of annual and perennial weeds and woody plants in forests, non-crop sites, and in and around aquatic sites; also for use in wildlife habitat areas, for perennial grass release, and grass growth suppression and grazed areas on these sites.

Avoid contact of herbicide with foliage, green stems, exposed non-woody roots or fruit of crops, desirable plants and trees, because severe injury or destruction may result.

### Active Ingredient(s):

|                                                                                  |        |
|----------------------------------------------------------------------------------|--------|
| glyphosate <sup>1</sup> N-(phosphonomethyl)glycine,<br>isopropylamine salt ..... | 53.8%  |
| Other Ingredients .....                                                          | 46.2%  |
| Total Ingredients .....                                                          | 100.0% |

<sup>1</sup> Contains 5.4 pounds per gallon glyphosate, isopropylamine salt (4 pounds per gallon glyphosate acid).

EPA Reg. No. 62719-324

## Keep Out of Reach of Children

## CAUTION PRECAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

#### Harmful If Inhaled

Avoid breathing spray mist. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

## Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE (Personal Protective Equipment). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

## First Aid

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 for emergency medical treatment information.

## Environmental Hazards

Do not contaminate water when cleaning equipment or disposing of equipment washwaters. Treatment of aquatic weeds can result in oxygen depletion or loss due to decomposition of dead plants. This oxygen loss can cause fish suffocation.

In case of leak or spill, soak up and remove to a landfill.

## Physical or Chemical Hazards

Spray solutions of this product should be mixed, stored and applied using only stainless steel, aluminum, fiberglass, plastic or plastic-lined steel containers.

Do not mix, store or apply this product or spray solutions of this product in galvanized steel or unlined steel (except stainless steel) containers or spray tanks. This product or spray solutions of this product react with such containers and tanks to produce hydrogen gas, which may form a highly combustible gas mixture. This gas mixture could flash or explode, causing serious personal injury, if ignited by open flame, spark, welder's torch, lighted cigarette or other ignition source.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Terms and Conditions of Use, Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at [www.dowagro.com](http://www.dowagro.com).

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

### Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

**This is an end-use product. Dow AgroSciences does not intend and has not registered it for reformulation. See individual container label for repackaging limitations.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

### Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

### Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep people and pets off treated areas until spray solution has dried.

### Storage and Disposal

Do not contaminate water, food, feed or seed by storage or disposal.

**Pesticide Storage:** Store above 10°F (-12°C) to keep product from crystallizing. Crystals will settle to the bottom. If allowed to crystallize, place in a warm room 68°F (20°C) for several days to redissolve and roll or shake container or recirculate in mini-bulk containers to mix well before using.

**Pesticide Disposal:** Wastes resulting from use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable Federal, state or local procedures.

### Nonrefillable containers 5 gallons or less:

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

### Refillable containers larger than 5 gallons:

**Container Reuse:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. If practical, agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

### Nonrefillable containers larger than 5 gallons:

**Container Reuse:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse** as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse** as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

## **General Information**

### **(How this product works)**

This product is a water-soluble liquid, which mixes readily with water and nonionic surfactant to be applied as a foliar spray for the control or destruction of many herbaceous and woody plants. This product is intended for control of annual and perennial weeds and woody plants in forests, pine straw plantations, non-crop sites such as utility rights-of-way, and in and around aquatic sites; also for use in wildlife habitat areas, for perennial grass release, and grass growth suppression and grazed areas on these sites.

The active ingredient in this product moves through the plant from the point of foliage contact to and into the root system. Visible effects on most annual weeds occur within 2 to 4 days, 7 days or more on most perennial weeds, and 30 days or more on most woody plants. Extremely cool or cloudy weather following treatment may slow the activity of this product and delay visual effects of control. Visible effects include gradual wilting and yellowing of the plant which advances to complete browning of above-ground growth and deterioration of underground plant parts.

Unless otherwise directed on this label, delay application until vegetation has emerged and reached the stages described for control of such vegetation under the "Weeds Controlled" section of this label.

Unemerged plants arising from unattached underground rhizomes or root stocks of perennials or brush will not be affected by the spray and will continue to grow. For this reason best control of most perennial weeds or brush is obtained when treatment is made at late growth stages approaching maturity.

Always use the higher rate of this product and surfactant within the recommended range when vegetation is heavy or dense, when treating dense multi-canopied sites or woody vegetation or difficult-to-control herbaceous or woody plants.

Do not treat weeds, brush or trees under poor growing conditions such as drought stress, disease or insect damage, as reduced control may result. Reduced control of target vegetation may also occur if foliage is heavily covered with dust at the time of treatment.

Reduced control may result when applications are made to woody plants or weeds following site disturbance or plant top growth removal from grazing, mowing, logging or mechanical brush control. For best results, delay treatment of such areas until resprouting and foliar growth has restored the target vegetation to the recommended stage of growth for optimum herbicide exposure and control.

Rainfall or irrigation occurring within 6 hours after application may reduce effectiveness. Heavy rainfall or irrigation within 2 hours after application may wash the product off the foliage and a repeat treatment may be required.

This product does not provide residual weed control. For subsequent residual weed control, follow a label-approved herbicide program. Read and carefully observe the cautionary statements and all other information appearing on the labels of all herbicides used.

**Note:** The maximum rates stated throughout this product's labeling apply to this product combined with the use of all other herbicides containing glyphosate or sulfosate as the active ingredient, whether applied as mixtures or separately. Calculate the application rates and ensure that the total use of this and other glyphosate or sulfosate containing products does not exceed the maximum use rates.

**Grazing Restrictions:** This product may be used to treat undesirable vegetation in utility rights-of-way that pass through pastures, rangeland, and forestry sites that are being grazed. For tank mix applications, comply with all restrictions appearing on the tank mix product label.

Except for lactating dairy animals there are no grazing restrictions following the labeled applications of this product.

- For lactating dairy animals there are no grazing restrictions for the following labeled applications of this product:
  - ▶ Where the spray can be directed onto undesirable woody brush and trees, such as in handgun spray-to-wet or low volume directed spray treatments.
  - ▶ For tree injection of frill applications and for cut stump treatments
- For broadcast applications, observe the following restrictions for lactating dairy animals:
  - ▶ For application rates of greater than 4.5 but not to exceed 7.5 quarts per acre, no more than 15 percent of the available grazing area may be treated.
  - ▶ For application rates that do not exceed 4.5 quarts per acre, no more than 25 percent of the available grazing area may be treated.
- These restrictions do not apply to pastures, rangeland or forestry sites outside of utility rights-of-way.

**NOTE:** Use of this product in any manner not consistent with this label may result in injury to persons, animals or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

Buyer and all users are responsible for all loss or damage in connection with the use or handling of mixtures of this product or other materials that are not expressly recommended in this label. Mixing this product with herbicides or other materials not recommended in this label may result in reduced performance.

**ATTENTION:** Avoid drift. Extreme care must be used when applying this product to prevent injury to desirable plants and crops.

Do not allow the herbicide solution to mist, drip, drift or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to the crop, plants or other areas on which treatment was not intended. The likelihood of plant or crop injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **Avoid applying at excessive speed or pressure.**

## Spray Drift Management

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees. Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following **Aerial Drift Reduction Advisory Information:**

**Importance of Droplet Size:** The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion section of this label).

**Controlling Droplet Size:** Volume-Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows product larger droplets.

Pressure-Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles-Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation-Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type-Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length-For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

Application-Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

**Swath Adjustment:** When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

**Wind:** Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

**Temperature and Humidity:** When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

**Temperature Inversions:** Applications should not occur during a temperature inversion, because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a connected cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

**Sensitive Areas:** The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

---

## Mixing And Application Instructions

---

Apply these spray solutions in properly maintained and calibrated equipment capable of delivering desired volumes. Hand-gun applications should be properly directed to avoid spraying desirable plants. Note: reduced results may occur if water containing soil is used, such as water from ponds and unlined ditches.

### Mixing

This product mixes readily with water. Mix spray solutions of this product as follows:

1. Fill the mixing or spray tank with the required amount of water while adding the required amount of this product (see "Directions for Use" and "Weeds Controlled" sections of this label).
2. Near the end of the filling process, add the required surfactant and mix well. Remove hose from tank immediately after filling to avoid siphoning back into the water source.

**Note:** If tank mixing with Garlon® 3A herbicide, ensure that Garlon 3A is well mixed with at least 75 percent of the total spray volume before adding this product to the spray tank to avoid incompatibility.

During mixing and application, foaming of the spray solution may occur. To prevent or minimize foam, avoid the use of mechanical agitators, place the filling hose below the surface of the spray solution (only during filling), terminate by-pass and return lines at the bottom of the tank, and, if needed, use an approved anti-foam or defoaming agent.

Keep by-pass line on or near bottom of tank to minimize foaming. Screen size in nozzle or line strainers should be no finer than 50 mesh. Carefully select correct nozzle to avoid spraying a fine mist. For best results with conventional ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**IMPORTANT:** When using this product, unless otherwise specified, mix with a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. For conifer release (pine release) use only surfactants that are approved for conifer release, and specified on the surfactant label as safe for use in conifer release (pine release). Always read and follow the manufacturer's surfactant label recommendations for best results.

Colorants or marking dyes approved for use with herbicides may be added to spray mixtures of this product. Colorants or dyes used in spray solutions of this product may reduce performance, especially at lower rates or dilutions. Use colorants or dyes according to the manufacturer's label recommendations.

Clean sprayer and parts immediately after using this product by thoroughly flushing with water and dispose of rinsate according to labeled use or disposal instructions.

Carefully observe all cautionary statements and other information appearing in the surfactant label.

### **Application Equipment And Techniques**

**ATTENTION: AVOID DRIFT. EXTREME CARE MUST BE EXERCISED WHEN APPLYING THIS PRODUCT TO PREVENT INJURY TO DESIRABLE PLANTS AND CROPS.**

Do not allow the herbicide solution to mist, drip, drift, or splash onto desirable vegetation since minute quantities of this product can cause severe damage or destruction to crops, plants, or other areas on which the treatment was not intended. The likelihood of plant or crop injury occurring from the use of this product is greatest when winds are gusty or in excess of 5 miles per hour or when other conditions, including lesser wind velocities, will allow spray drift to occur. When spraying, avoid combinations of pressure and nozzle type that will result in splatter or fine particles (mist) which are likely to drift. **AVOID APPLYING AT EXCESSIVE SPEED OR PRESSURE.**

**Note:** Use of this product in a manner not consistent with this label may result in injury to persons, animals, or crops, or other unintended consequences. When not in use, keep container closed to prevent spills and contamination.

### **Aerial Equipment**

**For aerial application of this product in California, refer to Federal supplemental label for this product entitled "For Aerial Application in California Only".** In California, aerial application may be made in aquatic sites and noncrop areas, including aquatic sites present in noncrop areas that are part of the intended treatment.

**For control of weed or brush species listed in this label using aerial application equipment:** For aerial broadcast application, unless otherwise specified, apply the rates of this product and surfactant recommended for broadcast application in a spray volume of 3 to 20 gallons of water per acre. See the "Weeds Controlled" section of this label for labeled annual and herbaceous weeds and woody plants and broadcast rate recommendations. Aerial applications of this product may only be made as specifically recommended in this label.

**AVOID DRIFT. Do not apply during inversion conditions, when winds are gusty or under any other condition which will allow drift. Drift may cause damage to any vegetation contacted to which treatment is not intended. To prevent injury to adjacent desirable vegetation, appropriate buffer zones must be maintained.**

Coarse sprays are less likely to drift; therefore, do not use nozzles or nozzle configurations which dispense spray as fine spray droplets. Do not angle nozzles forward into the airstream and do not increase spray volume by increasing nozzle pressure.

Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing in the additive label. The use of a drift control agent for conifer and herbaceous release applications may result in conifer injury and is not recommended.

**Ensure uniform application.** To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Thoroughly wash aircraft, especially landing gear, after each day of spraying to remove residues of this product accumulated during spraying or from spills. **Prolonged exposure of this product to uncoated steel surfaces may result in corrosion and possible failure of the part. Landing gear are most susceptible.** The maintenance of an organic coating (paint) which meets aerospace specification MIL-C-38413 may prevent corrosion.

### **Ground Broadcast Equipment**

**For control of weed or brush species listed in this label using conventional boom equipment:** For ground broadcast application, unless otherwise specified, apply the rates of this product and surfactant recommended for broadcast application in a spray volume of 3 to 30 gallons of water per acre. See the "Weeds Controlled" section of this label for labeled annual and herbaceous weeds and woody plants and broadcast rate recommendations. As density of vegetation increases, spray volume should be increased within the recommended range to ensure complete coverage. Carefully select correct nozzle to avoid spraying a fine mist. For best results with ground application equipment, use flat fan nozzles. Check for even distribution of spray droplets.

**Forestry and Utility Rights-of-Way Sites:** This product is recommended for broadcast applications using suitable ground equipment in forestry sites, utility sites, and utility rights-of-way. Apply the recommended rates of this product and surfactant in a spray volume of 10 to 60 gallons per acre. Check for even distribution of spray droplets.

### Hand-Held and High-Volume Equipment (Use Coarse Sprays Only)

**For control of weeds listed in this label using knapsack sprayers or high-volume spraying equipment utilizing handguns or other suitable nozzle arrangements:**

**High volume sprays:** Prepare a 3/4 to 2 percent solution of this product in water, add a nonionic surfactant and apply to foliage of vegetation to be controlled. For specific rates of application and instructions for control of various annual and perennial weeds, see the "Weeds Controlled" section in this label.

Prepare the desired volume of spray solution by mixing the amount of this product in water, shown in the following table:

#### Spray Solution

| Desired Volume | Amount of this product |             |             |           |             |             |              |              |
|----------------|------------------------|-------------|-------------|-----------|-------------|-------------|--------------|--------------|
|                | 3/4%                   | 1%          | 1 1/4%      | 1 1/2%    | 2%          | 5%          | 8%           | 10%          |
| 1 gal          | 1 fl oz                | 1 1/3 fl oz | 1 2/3 fl oz | 2 fl oz   | 2 2/3 fl oz | 6 1/2 fl oz | 10 1/4 fl oz | 12 3/4 fl oz |
| 25 gal         | 1 1/2 pt               | 1 qt        | 1 1/4 qt    | 1 1/2 qt  | 2 qt        | 5 qt        | 2 gal        | 2.5 gal      |
| 100 gal        | 3 qt                   | 1 gal       | 1 1/4 gal   | 1 1/2 gal | 2 gal       | 5 gal       | 8 gal        | 10 gal       |

2 tablespoons = 1 fluid ounce

For use in knapsack sprayers, it is suggested that the recommended amount of this product be mixed with water in a larger container. Fill the knapsack sprayer with the mixed solution and add the correct amount of surfactant.

### Selective Equipment

This product may be applied through shielded sprayers or wiper application equipment. This equipment may be used to selectively control undesirable vegetation without harming desirable vegetation.

Shielded sprayers direct the herbicide solution onto weeds while shielding desirable vegetation from the spray solution. Any recommended rate or tank mixture of this product may be used employing this equipment.

Wiper applicators physically wipe product directly onto undesirable vegetation. Care should be taken to avoid wiping desirable vegetation. Use a 33 to 100 percent solution of this product, diluted in water for wiper applications. Use a 33 percent solution for wick or gravity feed systems. Higher concentrations may be used in pressurized systems that are capable of handling thicker solutions. Addition of a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution is recommended.

Applications should be made on a spray-to-wet basis. Spray coverage should be uniform and complete. Do not spray to point of runoff.

**Low volume directed sprays:** This product may be used as a 5 to 10 percent solution in low-volume directed sprays for spot treatment of trees and brush. This treatment method is most effective in areas where there is a low density of undesirable trees or brush. If a straight stream nozzle is used, start the application at the top of the targeted vegetation and spray from top to bottom in a lateral zig-zag motion. Ensure that at least 50 percent of the leaves are contacted by the spray solution. For flat fan and cone nozzles and with hand-directed mist blowers, mist the application over the foliage of the targeted vegetation. Small, open-branched trees need only be treated from one side. If the foliage is thick or there are multiple root sprouts, applications must be made from several sides to ensure adequate spray coverage.

## Weeds Controlled

### Annual Weeds

Apply to actively growing annual grasses and broadleaf weeds.

Allow at least 3 days after application before disturbing treated vegetation. After this period the weeds may be mowed, tilled or burned. See "Directions for Use," "General Information" and "Mixing and Application Instructions" for labeled uses and specific application instructions.

**Broadcast Application Rates:** For weeds less than 6 inches tall, use 1 1/2 pints of this product per acre plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. If weeds are greater than 6 inches tall, use 2 1/2 pints of this product per acre plus a non-ionic surfactant containing 80% or greater active ingredient.

**Hand-Held, High-Volume Application Rates:** Use a 3/4 percent solution of this product in water plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Apply to foliage of vegetation to be controlled.

When applied as directed, this product plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient will control the following annual weeds:

| Common Name                     | Scientific Name                 |
|---------------------------------|---------------------------------|
| Balsamapple <sup>†</sup>        | <i>Momordica charantia</i>      |
| Barley                          | <i>Hordeum vulgare</i>          |
| Barnyardgrass                   | <i>Echinochloa crus-galli</i>   |
| Bassia, fivehook                | <i>Bassia hyssopifolia</i>      |
| Bluegrass, annual               | <i>Poa annua</i>                |
| Bluegrass, bulbous              | <i>Poa bulbosa</i>              |
| Brome                           | <i>Bromus spp.</i>              |
| Buttercup                       | <i>Ranunculus spp.</i>          |
| Cheat                           | <i>Bromus secalinus</i>         |
| Chickweed, mouseear             | <i>Cerastium vulgatum</i>       |
| Cocklebur                       | <i>Xanthium strumarium</i>      |
| Corn, volunteer                 | <i>Zea mays</i>                 |
| Crabgrass                       | <i>Digitaria spp.</i>           |
| Dwarf dandelion                 | <i>Krigia cespitosa</i>         |
| Falseflax, smallseed            | <i>Camelina microcarpa</i>      |
| Fiddleneck                      | <i>Amsinckia spp.</i>           |
| Flaxleaf fleabane               | <i>Conyza bonariensis</i>       |
| Fleabane                        | <i>Erigeron spp.</i>            |
| Foxtail                         | <i>Setaria spp.</i>             |
| Foxtail, Carolina               | <i>Alopecurus carolinianus</i>  |
| Groundsel, common               | <i>Senecio vulgaris</i>         |
| Horseweed/Marestail             | <i>Conyza canadensis</i>        |
| Kochia                          | <i>Kochia scoparia</i>          |
| Lambsquarters, common           | <i>Chenopodium album</i>        |
| Lettuce, prickly                | <i>Lactuca serriola</i>         |
| Morningglory                    | <i>Ipomoea spp.</i>             |
| Mustard, blue                   | <i>Chorispora tenella</i>       |
| Mustard, tansy                  | <i>Descurainia pinnata</i>      |
| Mustard, tumble                 | <i>Sisymbrium altissimum</i>    |
| Mustard, wild                   | <i>Sinapis arvensis</i>         |
| Oats, wild                      | <i>Avena fatua</i>              |
| Panicum                         | <i>Panicum spp.</i>             |
| Pennycress, field               | <i>Thlaspi arvense</i>          |
| Pigweed, redroot                | <i>Amaranthus retroflexus</i>   |
| Pigweed, smooth                 | <i>Amaranthus hybridus</i>      |
| Ragweed, common                 | <i>Ambrosia artemisiifolia</i>  |
| Ragweed, giant                  | <i>Ambrosia trifida</i>         |
| Rocket, London                  | <i>Sisymbrium irio</i>          |
| Rye                             | <i>Secale cereale</i>           |
| Ryegrass, Italian <sup>††</sup> | <i>Lolium multiflorum</i>       |
| Sandbur, field                  | <i>Cenchrus spp.</i>            |
| Shattercane                     | <i>Sorghum bicolor</i>          |
| Shepherd's-purse                | <i>Capsella bursa-pastoris</i>  |
| Signalgrass, broadleaf          | <i>Brachiaria platyphylla</i>   |
| Smartweed, Pennsylvania         | <i>Polygonum pennsylvanicum</i> |
| Sowthistle, annual              | <i>Sonchus oleraceus</i>        |
| Spanishneedles <sup>††</sup>    | <i>Bidens bipinnata</i>         |
| Stinkgrass                      | <i>Eragrostis cilianensis</i>   |
| Sunflower                       | <i>Helianthus annuus</i>        |
| Thistle, Russian                | <i>Salsola kali</i>             |
| Spurry, umbrella                | <i>Holosteum umbellatum</i>     |
| Velvetleaf                      | <i>Abutilon theophrasti</i>     |
| Wheat                           | <i>Triticum aestivum</i>        |
| Witchgrass                      | <i>Panicum capillare</i>        |

<sup>†</sup> Apply with hand-held equipment only.

<sup>††</sup> Apply 3 pints of this product per acre.

Annual weeds will generally continue to germinate from seed throughout the growing season. Repeat treatments will be necessary to control later germinating weeds.

## Perennial Weeds

Apply this product to control most vigorously growing perennial weeds. Unless otherwise directed, apply when target plants are actively growing and most have reached early head or early bud stage of growth. Unless otherwise directed, allow at least 7 days after application before disturbing vegetation.

**NOTE:** If weeds have been mowed or tilled, do not treat until regrowth has reached the recommended stages. Fall treatments must be applied before a killing frost.

Repeat treatments may be necessary to control weeds regenerating from underground parts or seed.

**Specific Weed Control Recommendations:** For perennial weeds, apply the recommended rate plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

**When applied as directed, this product plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient will control the following perennial weeds:** (Numbers in parentheses "(-)" following common name of a listed weed species refer to "Specific Perennial Weed Control Recommendations" for that weed which follow the species listing.)

| Common Name                      | Scientific Name                      |
|----------------------------------|--------------------------------------|
| Alfalfa (31)                     | <i>Medicago sativa</i>               |
| Alligatorweed <sup>†</sup> (1)   | <i>Alternanthera philoxeroides</i>   |
| Anise/Fennel (31)                | <i>Foeniculum vulgare</i>            |
| Artichoke, Jerusalem (31)        | <i>Helianthus tuberosus</i>          |
| Bahiagrass (31)                  | <i>Paspalum notatum</i>              |
| Bermudagrass (2)                 | <i>Cynodon dactylon</i>              |
| Bindweed, field (3)              | <i>Convolvulus arvensis</i>          |
| Bluegrass, Kentucky (12)         | <i>Poa pratensis</i>                 |
| Blueweed, Texas (3)              | <i>Helianthus ciliaris</i>           |
| Brackenfern (4)                  | <i>Pteridium spp.</i>                |
| Bromegrass, smooth (12)          | <i>Bromus inermis</i>                |
| Canarygrass, reed (12)           | <i>Phalaris arundinacea</i>          |
| Cattail (5)                      | <i>Typha spp.</i>                    |
| Clover, red (31)                 | <i>Trifolium pratense</i>            |
| Clover, white (31)               | <i>Trifolium repens</i>              |
| Cogongrass (6)                   | <i>Imperata cylindrica</i>           |
| Cordgrass (7)                    | <i>Spartina spp.</i>                 |
| Cutgrass, giant <sup>†</sup> (8) | <i>Zizaniopsis miliacea</i>          |
| Dallisgrass (31)                 | <i>Paspalum dilatatum</i>            |
| Dandelion (31)                   | <i>Taraxacum officinale</i>          |
| Dock, curly (31)                 | <i>Rumex crispus</i>                 |
| Dogbane, hemp (9)                | <i>Apocynum cannabinum</i>           |
| Fescue (31)                      | <i>Festuca spp.</i>                  |
| Fescue, tall (10)                | <i>Festuca arundinacea</i>           |
| Guineagrass (11)                 | <i>Panicum maximum</i>               |
| Hemlock, poison (31)             | <i>Conium maculatum</i>              |
| Horsenettle (31)                 | <i>Solanum carolinense</i>           |
| Horseradish (9)                  | <i>Armoracia rusticana</i>           |
| Ice Plant (22)                   | <i>Mesembryanthemum crystallinum</i> |
| Johnsongrass (12)                | <i>Sorghum halepense</i>             |
| Kikuyugrass (21)                 | <i>Pennisetum clandestinum</i>       |

| Common Name                          | Scientific Name               |
|--------------------------------------|-------------------------------|
| Knapweed (9)                         | <i>Centaurea repens</i>       |
| Lantana (13)                         | <i>Lantana camara</i>         |
| Lespedeza, common (31)               | <i>Lespedeza striata</i>      |
| Lespedeza, sericea (31)              | <i>Lespedeza cuneata</i>      |
| Loosestrife, purple (14)             | <i>Lythrum salicaria</i>      |
| Lotus, American (15)                 | <i>Nelumbo lutea</i>          |
| Maidencane (16)                      | <i>Panicum hematomon</i>      |
| Milkweed (17)                        | <i>Asclepias spp.</i>         |
| Muhly, wirestem (21)                 | <i>Muhlenbergia frondosa</i>  |
| Mullein, common (31)                 | <i>Verbascum thapsus</i>      |
| Napiergrass (31)                     | <i>Pennisetum purpureum</i>   |
| Nightshade, silverleaf (3)           | <i>Solanum elaeagnifolium</i> |
| Nutsedge, purple (18)                | <i>Cyperus rotundus</i>       |
| Nutsedge, yellow (18)                | <i>Cyperus esculentus</i>     |
| Orchardgrass (12)                    | <i>Dactylis glomerata</i>     |
| Pampasgrass (19)                     | <i>Cortaderia jubata</i>      |
| Paragrass (16)                       | <i>Brachiaria mutica</i>      |
| Phragmites <sup>††</sup> (20)        | <i>Phragmites spp.</i>        |
| Quackgrass (21)                      | <i>Agropyron repens</i>       |
| Reed, giant (22)                     | <i>Arundo donax</i>           |
| Ryegrass, perennial (12)             | <i>Lolium perenne</i>         |
| Smartweed, swamp (31)                | <i>Polygonum coccineum</i>    |
| Spatterdock (23)                     | <i>Nuphar luteum</i>          |
| Starthistle, yellow (31)             | <i>Centaurea solstitialis</i> |
| Sweet potato, wild <sup>†</sup> (24) | <i>Ipomoea pandurata</i>      |
| Thistle, artichoke (25)              | <i>Cynara cardunculus</i>     |
| Thistle, Canada (25)                 | <i>Cirsium arvense</i>        |
| Timothy (12)                         | <i>Phleum pratense</i>        |
| Torpedograss <sup>†</sup> (26)       | <i>Panicum repens</i>         |
| Tules, common (27)                   | <i>Scirpus acutus</i>         |
| Vaseygrass (31)                      | <i>Paspalum urvillei</i>      |
| Velvetgrass (31)                     | <i>Holcus spp.</i>            |
| Waterhyacinth (28)                   | <i>Eichornia crassipes</i>    |
| Waterlettuce (29)                    | <i>Pistia stratiotes</i>      |
| Waterprimrose (30)                   | <i>Ludwigia spp.</i>          |
| Wheatgrass, western (12)             | <i>Agropyron smithii</i>      |

<sup>†</sup> Partial control.

<sup>††</sup> Partial control in southeastern states. See "Specific Weed Control Recommendations" below.

#### Specific Perennial Weed Control Recommendations:

- Alligatorweed:** Apply 6 pints of this product per acre as a broadcast spray or as a 1 1/4 percent solution with hand-held equipment to provide partial control of alligatorweed. Apply when most of the target plants are in bloom. Repeat applications will be required to maintain such control.
- Bermudagrass:** Apply 7 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and when seedheads appear.
- Blindweed, field / Silverleaf Nightshade / Texas Blueweed:** Apply 6 to 7 1/2 pints of this product per acre as a broadcast spray west of the Mississippi River and 4 1/2 to 6 pints of this product per acre east of the Mississippi River. With hand-held equipment, use a 1 1/2 percent solution. Apply when target plants are actively growing and are at or beyond full bloom. For silverleaf nightshade, best results can be obtained when application is made after berries are formed. Do not treat when weeds are under drought stress. New leaf development indicates active growth. For best results apply in late summer or fall.
- Brackenfern:** Apply 4 1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 to 1 percent solution with hand-held equipment. Apply to fully expanded fronds which are at least 18 inches long.
- Cattail:** Apply 4 1/2 to 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and are at or beyond the early-to-full bloom stage of growth. Best results are achieved when application is made during the summer or fall months.
- Cogongrass:** Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray. Apply when cogongrass is at least 18 inches tall and actively growing in late summer or fall. Allow 7 or more days after application before tillage or mowing. Due to uneven stages of growth and the dense nature of vegetation preventing good spray coverage, repeat treatments may be necessary to maintain control.
- Cordgrass:** Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 to 2 percent solution with hand-held equipment. Schedule applications in order to allow 6 hours before treated plants are covered by tidewater. The presence of debris and silt on the cordgrass plants will reduce performance. It may be necessary to wash targeted plants prior to application to improve uptake of this product into the plant.
- Cutgrass, giant:** Apply 6 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment to provide partial control of giant cutgrass. Repeat applications will be required to maintain such control, especially where vegetation is partially submerged in water. Allow for substantial regrowth to the 7 to 10-leaf stage prior to retreatment.
- Dogbane, hemp / Knapweed / Horseradish:** Apply 6 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the late bud-to-flower stage of growth. For best results, apply in late summer or fall.
- Fescue, tall:** Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 1 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained.
- Gulmeagrass:** Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and when most have reached at least the 7-leaf stage of growth.
- Johnsongrass / Bluegrass, Kentucky / Bromegrass, smooth / Canarygrass, reed / Orchardgrass / Ryegrass, perennial / Timothy / Wheatgrass, western:** Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the boot-to-head stage of growth. When applied prior to the boot stage, less desirable control may be obtained. In the fall, apply before plants have turned brown.
- Lantana:** Apply this product as a 3/4 to 1 percent solution with hand-held equipment. Apply to actively growing lantana at or beyond the bloom stage of growth. Use the higher application rate for plants that have reached the woody stage of growth.
- Loosestrife, purple:** Apply 4 pints of this product per acre as a broadcast spray or as a 1 to 1 1/2 percent solution using hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost.

15. **Lotus, American:** Apply 4 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Treat when plants are actively growing at or beyond the bloom stage of growth. Best results are achieved when application is made during summer or fall months. Fall treatments must be applied before a killing frost. Repeat treatment may be necessary to control regrowth from underground parts and seeds.
16. **Maldenecane / Paragrass:** Apply 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Repeat treatments will be required, especially to vegetation partially submerged in water. Under these conditions, allow for regrowth to the 7 to 10-leaf stage prior to retreatment.
17. **Milkweed, common:** Apply 4 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Apply when target plants are actively growing and most have reached the late bud-to-flower stage of growth.
18. **Nutsedge, purple, yellow:** Apply 4 1/2 pints of this product per acre as a broadcast spray, or as a 3/4 percent solution with hand-held equipment to control existing nutsedge plants and immature nutlets attached to treated plants. Apply when target plants are in flower or when new nutlets can be found at rhizome tips. Nutlets which have not germinated will not be controlled and may germinate following treatment. Repeat treatments will be required for long-term control.
19. **Pampasgrass:** Apply a 1 1/2 percent solution of this product with hand-held equipment when plants are actively growing.
20. **Phragmites:** For partial control of phragmites in Florida and the counties of other states bordering the Gulf of Mexico, apply 7 1/2 pints per acre as a broadcast spray or apply a 1 1/2 percent solution with hand-held equipment. In other areas of the U.S., apply 4 to 6 pints per acre as a broadcast spray or apply a 3/4 percent solution with hand-held equipment for partial control. For best results, treat during late summer or fall months when plants are actively growing and in full bloom. Due to the dense nature of the vegetation, which may prevent good spray coverage and uneven stages of growth, repeat treatments may be necessary to maintain control. Visual control symptoms will be slow to develop.
21. **Quackgrass / Kikuyugrass / Muhly, wirestem:** Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment when most quackgrass or wirestem muhly is at least 8 inches in height (3 to 4-leaf stage of growth) and actively growing. Allow 3 or more days after application before tillage.
22. **Reed, giant / ice plant:** For control of giant reed and ice plant, apply a 1 1/2 percent solution of this product with hand-held equipment when plants are actively growing. For giant reed, best results are obtained when applications are made in late summer to fall.
23. **Spatterdock:** Apply 6 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Apply when most plants are in full bloom. For best results, apply during the summer or fall months.
24. **Sweet potato, wild:** Apply this product as a 1 1/2 percent solution using hand-held equipment. Apply to actively growing weeds that are at or beyond the bloom stage of growth. Repeat applications will be required. Allow the plant to reach the recommended stage of growth before retreatment.
25. **Thistle, Canada / artichoke:** Apply 3 to 4 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment for Canada thistle. To control artichoke thistle, apply a 2 percent solution as a spray-to-wet application. Apply when target plants are actively growing and are at or beyond the bud stage of growth.
26. **Torpedograss:** Apply 6 to 7 1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/2 percent solution with hand-held equipment to provide partial control of torpedograss. Use the lower rates under terrestrial conditions, and the higher rates under partially submerged or a floating mat condition. Repeat treatments will be required to maintain such control.
27. **Tules, common:** Apply this product as a 1 1/2 percent solution with hand-held equipment. Apply to actively growing plants at or beyond the seedhead stage of growth. After application, visual symptoms will be slow to appear and may not occur for 3 or more weeks.
28. **Waterhyacinth:** Apply 5 to 6 pints of this product per acre as a broadcast spray or apply a 3/4 to 1 percent solution with hand-held equipment. Apply when target plants are actively growing and at or beyond the early bloom stage of growth. After application, visual symptoms may require 3 or more weeks to appear with complete necrosis and decomposition usually occurring within 60 to 90 days. Use the higher rates when more rapid visual effects are desired.
29. **Waterlettuce:** For control, apply a 3/4 to 1 percent solution of this product with hand-held equipment to actively growing plants. Use higher rates where infestations are heavy. Best results are obtained from mid-summer through winter applications. Spring applications may require retreatment.
30. **Waterprimrose:** Apply this product as a 3/4 percent solution using hand-held equipment. Apply to plants that are actively growing at or beyond the bloom stage of growth, but before fall color changes occur. Thorough coverage is necessary for best control.
31. **Other perennial weeds listed above:** Apply 4 1/2 to 7 1/2 pints of this product per acre as a broadcast spray or apply as a 3/4 to 1 1/2 percent solution with hand-held equipment.

## Woody Brush and Trees

**NOTE:** If brush has been mowed or tilled or trees have been cut, do not treat until regrowth has reached the recommended stage of growth.

### Application Rates and Timing

When applied as a 5 to 8 percent solution as a directed application as described in the "Hand-Held and High-Volume Equipment" section, this product will control or partially control all wood brush and tree species listed in this section of this label. Use the higher rate of application for dense stands and larger woody brush and trees.

**Specific Brush or Tree Control Recommendations:** Numbers in parentheses "(-)" following the common name of a listed brush or tree species refer to "Specific Brush or Tree Control Recommendations" which follow the species listing. See this section for specific application rates and timing for listed species.

For woody brush and trees, apply the recommended rate plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information. Make applications when plants are actively growing and, unless otherwise directed, after full-leaf expansion. Use the higher rate for larger plants and/or dense areas of growth. On vines, use the higher rate for plants that have reached the woody stage of growth. Best results are obtained when application is made in late summer or fall after fruit formation.

In arid areas, best results are obtained when application is made in the spring or early summer when brush species are at high moisture content and are flowering. Ensure thorough coverage when using hand-held equipment. Symptoms may not appear prior to frost or senescence with fall treatments.

Allow 7 or more days after application before tillage, mowing or removal. Repeat treatments may be necessary to control plants regenerating from underground parts or seed. Some autumn colors on undesirable deciduous species are acceptable provided no major leaf drop has occurred. Reduced performance may result if fall treatments are made following a frost.

See the "Directions for Use" and "Mixing and Application Instructions" sections in this label for labeled use and specific application instructions. **When applied as directed, this product plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient will control the following woody brush plants and trees:** (Numbers in parentheses "(-)" following common name of a listed brush or tree species refer to "Specific Brush or Tree Control Recommendations" for that species which follow the species listing.)

| Common Name                            | Scientific Name                    |
|----------------------------------------|------------------------------------|
| Alder (1)                              | <i>Alnus spp.</i>                  |
| Ash <sup>†</sup> (20)                  | <i>Fraxinus spp.</i>               |
| Aspen, quaking (2)                     | <i>Populus tremuloides</i>         |
| Bearclover, Bearmat (20)               | <i>Chamaebatia foliolosa</i>       |
| Birch (3)                              | <i>Betula spp.</i>                 |
| Blackberry (1)                         | <i>Rubus spp.</i>                  |
| Broom, French (4)                      | <i>Cytisus monspessulanus</i>      |
| Broom, Scotch (4)                      | <i>Cytisus scoparius</i>           |
| Buckwheat, California <sup>†</sup> (5) | <i>Eriogonum fasciculatum</i>      |
| Cascara <sup>†</sup> (20)              | <i>Rhamnus purshiana</i>           |
| Catsclaw <sup>†</sup> (6)              | <i>Acacia greggi</i>               |
| Ceanothus (20)                         | <i>Ceanothus spp.</i>              |
| Chamise (17)                           | <i>Adenostoma fasciculatum</i>     |
| Cherry, bitter (7)                     | <i>Prunus emarginata</i>           |
| Cherry, black (7)                      | <i>Prunus serotina</i>             |
| Cherry, pin (7)                        | <i>Prunus pensylvanica</i>         |
| Coyote brush (8)                       | <i>Baccharis consanguinea</i>      |
| Creeper, Virginia <sup>†</sup> (20)    | <i>Parthenocissus quinquefolia</i> |
| Dewberry (1)                           | <i>Rubus trivialis</i>             |
| Dogwood (9)                            | <i>Cornus spp.</i>                 |
| Elderberry (3)                         | <i>Sambucus spp.</i>               |
| Elm <sup>†</sup> (20)                  | <i>Ulmus spp.</i>                  |
| Eucalyptus, bluegum (10)               | <i>Eucalyptus globulus</i>         |
| Hasardia <sup>†</sup> (5)              | <i>Haplopappus squamosus</i>       |
| Hawthorn (2)                           | <i>Crataegus spp.</i>              |
| Hazel (3)                              | <i>Corylus spp.</i>                |
| Hickory (9)                            | <i>Carya spp.</i>                  |
| Holly, Florida (11)                    | <i>Schinus terebinthifolius</i>    |
| (Brazilian peppertree)                 |                                    |
| Honeysuckle (1)                        | <i>Lonicera spp.</i>               |
| Hornbeam, American (20)                | <i>Carpinus caroliniana</i>        |
| Kudzu (12)                             | <i>Pueraria lobata</i>             |
| Locust, black <sup>†</sup> (20)        | <i>Robinia pseudoacacia</i>        |
| Manzanita (20)                         | <i>Arctostaphylos spp.</i>         |
| Maple, red <sup>†</sup> (13)           | <i>Acer rubrum</i>                 |
| Maple, sugar (14)                      | <i>Acer saccharum</i>              |
| Maple, vine <sup>†</sup> (20)          | <i>Acer circinatum</i>             |
| Monkey flower <sup>†</sup> (5)         | <i>Mimulus guttatus</i>            |
| Oak, black <sup>†</sup> (20)           | <i>Quercus velutina</i>            |
| Oak, northern pin (14)                 | <i>Quercus palustris</i>           |
| Oak, post (1)                          | <i>Quercus stellata</i>            |

| Common Name                           | Scientific Name                |
|---------------------------------------|--------------------------------|
| Oak, red (14)                         | <i>Quercus rubra</i>           |
| Oak, southern red (7)                 | <i>Quercus falcata</i>         |
| Oak, white <sup>†</sup> (20)          | <i>Quercus alba</i>            |
| Persimmon <sup>†</sup> (20)           | <i>Diospyros spp.</i>          |
| Poison-ivy (15)                       | <i>Rhus radicans</i>           |
| Poison-oak (15)                       | <i>Rhus toxicodendron</i>      |
| Poplar, yellow <sup>†</sup> (20)      | <i>Liriodendron tulipifera</i> |
| Prunus (7)                            | <i>Prunus spp.</i>             |
| Raspberry (1)                         | <i>Rubus spp.</i>              |
| Redbud, eastern (20)                  | <i>Cercis canadensis</i>       |
| Rose, multiflora (16)                 | <i>Rosa multiflora</i>         |
| Russian-olive (20)                    | <i>Elaeagnus angustifolia</i>  |
| Sage: black (17), white               | <i>Salvia spp.</i>             |
| Sagebrush, California (17)            | <i>Artemisia californica</i>   |
| Salmonberry (3)                       | <i>Rubus spectabilis</i>       |
| Salt cedar <sup>†</sup> (9)           | <i>Tamarix spp.</i>            |
| Saltbush, sea myrtle (18)             | <i>Baccharis halimifolia</i>   |
| Sassafras (20)                        | <i>Sassafras albidum</i>       |
| Sourwood <sup>†</sup> (20)            | <i>Oxydendrum arboreum</i>     |
| Sumac, poison <sup>†</sup> (20)       | <i>Rhus vernix</i>             |
| Sumac, smooth <sup>†</sup> (20)       | <i>Rhus glabra</i>             |
| Sumac, winged <sup>†</sup> (20)       | <i>Rhus copallina</i>          |
| Sweetgum (7)                          | <i>Liquidambar styraciflua</i> |
| Swordfern <sup>†</sup> (20)           | <i>Polystichum munium</i>      |
| Tallowtree, Chinese (17)              | <i>Sapium sebiferum</i>        |
| Thimbleberry (3)                      | <i>Rubus parviflorus</i>       |
| Tobacco, tree <sup>†</sup> (5)        | <i>Nicotiana glauca</i>        |
| Trumpet creeper (2)                   | <i>Campsis radicans</i>        |
| Waxmyrtle, southern <sup>†</sup> (11) | <i>Myrica cerifera</i>         |
| Willow (19)                           | <i>Salix spp.</i>              |

<sup>†</sup> Partial control (See below for control or partial control instructions.)

#### Specific Brush or Tree Control Recommendations:

- Alder / Blackberry / Dewberry / Honeysuckle / Oak, Post / Raspberry:** For control, apply 4 1/2 to 6 pints per acre as a broadcast spray or as a 3/4 to 1 1/4 percent solution with hand-held equipment.
- Aspen, Quaking / Hawthorn / Trumpet creeper:** For control, apply 3 to 4 1/4 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/4 percent solution with hand-held equipment.
- Birch / Elderberry / Hazel / Salmonberry / Thimbleberry:** For control, apply 3 pints per acre of this product as a broadcast spray or as a 3/4 percent solution with hand-held equipment.
- Broom, French / Broom, Scotch:** For control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment.
- Buckwheat, California / Hasardia / Monkey flower / Tobacco, tree:** For partial control of these species, apply a 3/4 to 1 1/2 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
- Catsclaw:** For partial control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.
- Cherry, bitter / Cherry, black / Cherry, pin / Oak, southern red / Sweetgum / Prunus:** For control, apply 3 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 to 1 1/2 percent solution with hand-held equipment.
- Coyote brush:** For control, apply a 1 1/4 to 1 1/2 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.

9. **Dogwood / Hickory / Salt cedar:** For partial control, apply a 1 to 2 percent solution of this product with hand-held equipment or 6 to 7 1/2 pints per acre as a broadcast spray.
10. **Eucalyptus, bluegum:** For control of eucalyptus resprouts, apply a 1 1/2 percent solution of this product with hand-held equipment when resprouts are 6 to 12-feet tall. Ensure complete coverage. Apply when plants are actively growing. Avoid application to drought-stressed plants.
11. **Holly, Florida / Waxmyrtle, southern:** For partial control, apply this product as a 1 1/2 percent solution with hand-held equipment.
12. **Kudzu:** For control, apply 6 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Repeat applications will be required to maintain control.
13. **Maple, red:** For control, apply as a 3/4 to 1 1/4 percent solution with hand-held equipment when leaves are fully developed. For partial control, apply 2 to 7 1/2 pints of this product per acre as a broadcast spray.
14. **Maple, sugar / Oak: northern pin / Oak, red:** For control, apply as a 3/4 to 1 1/4 percent solution with hand-held equipment when at least 50 percent of the new leaves are fully developed.
15. **Poison-ivy / Poison-oak:** For control, apply 6 to 7 1/2 pints of this product per acre as a broadcast spray or as a 1 1/2 percent solution with hand-held equipment. Repeat applications may be required to maintain control. Fall treatments must be applied before leaves lose green color.
16. **Rose, multiflora:** For control, apply 3 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment. Treatments should be made prior to leaf deterioration by leaf-feeding insects.
17. **Sage, black / Sagebrush, California / Chamise / Tallowtree, Chinese:** For control of these species, apply a 3/4 percent solution of this product as a foliar spray with hand-held equipment. Thorough coverage of foliage is necessary for best results.
18. **Saltbush, sea myrtle:** For control, apply this product as a 1 percent solution with hand-held equipment.
19. **Willow:** For control, apply 4 1/2 pints of this product per acre as a broadcast spray or as a 3/4 percent solution with hand-held equipment.
20. **Other woody brush and trees listed above:** For partial control, apply 3 to 7 1/2 pints of this product per acre as a broadcast spray or as a 3/4 to 1 1/2 percent solution with hand-held equipment.

### **Aquatic and other Noncrop Sites**

Apply this product as directed and under conditions described to control or partially control weeds and woody plants listed in the "Weeds Controlled" section in industrial, recreational and public areas or other similar aquatic or terrestrial sites on this label.

#### **Noncrop Sites**

This product may be used to control the listed weeds in and around aquatic sites and on noncrop sites such as :

Airports  
Golf Courses  
Habitat Restoration & Management Areas  
Highways & Roadsides  
Industrial Plant Sites  
Lumberyards  
Parking Areas  
Parks  
Petroleum Tank Farms

Pipeline, Power, Telephone & Utility Rights-of-Way  
Pumping Installations  
Railroads  
Schools  
Storage Areas  
Similar Sites

#### **Aquatic Sites**

This product may be applied to emerged weeds in all bodies of fresh and brackish water which may be flowing, nonflowing or transient. This includes lakes, rivers, streams, ponds, estuaries, rice levees, seeps, irrigation and drainage ditches, canals, reservoirs, wastewater treatment facilities, wildlife habitat restoration and management areas and similar sites.

If aquatic sites are present in the noncrop area and are part of the intended treatment, read and observe the following directions:

- This product does not control plants which are completely submerged or have a majority of their foliage under water.
- There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.
- Consult local state fish and game agency and water control authorities before applying this product to public water. Permits may be required to treat such water.
- **NOTE:** Do not apply this product directly to water within 1/2 mile upstream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after the application. The water intake may be turned on prior to 48 hours if the glyphosate level in the intake water is below 0.7 parts per million as determined by laboratory analysis. These aquatic applications may be made **only** in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the applications. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.
- For treatments after drawdown of water or in dry ditches, allow 7 or more days after treatment before reintroduction of water to achieve maximum weed control. Apply this product within 1 day after drawdown to ensure application to actively growing weeds.
- Floating mats of vegetation may require retreatment. Avoid wash-off of sprayed foliage by spray boat or recreational boat backwash or by rainfall within 6 hours of application. Do not re-treat within 24 hours following the initial treatment.
- Applications made to moving bodies of water must be made while traveling upstream to prevent concentration of this herbicide in water. When making any bankside applications, do not overlap more than 1 foot into open water. Do not spray in bodies of water where weeds do not exist. The maximum application rate of 7 1/2 pints per acre must not be exceeded in any single broadcast application that is being made over water.

- When emerged infestations require treatment of the total surface area of impounded water, treating the area in strips may avoid oxygen depletion due to decaying vegetation. Oxygen depletion may result in fish kill.

### Forestry Sites and Utility Rights-of-Way

In forest and utility sites, this product is recommended for the control or partial control of woody brush, trees, and annual and perennial herbaceous weeds. This product is also recommended for use in preparing or establishing wildlife openings within these sites, in pine straw plantations for maintaining logging roads, and for side trimming along utility rights-of-way.

In forestry sites, this product is recommended for use in site preparation prior to planting any tree species, including Christmas trees and silvicultural nursery sites.

In utility sites, this product is recommended for use along electrical power, pipeline, and telephone rights-of-way, and in other utility sites associated with these rights-of-way, such as substations.

#### Application Rates <sup>1</sup>:

| Method of Application                         | Application Rate     | Spray Volume (gal/acre) |
|-----------------------------------------------|----------------------|-------------------------|
| <b>Broadcast</b>                              |                      |                         |
| Aerial                                        | 1.5 to 7.5 qt/acre   | 5 to 30                 |
| Ground                                        | 1.5 to 7.5 qt/acre   | 10 to 60                |
| <b>Spray-to-Wet</b>                           |                      |                         |
| Handgun, Backpack Mistblower                  | 0.75 to 2% by volume | spray-to-wet            |
| <b>Low Volume Directed Spray <sup>2</sup></b> |                      |                         |
| Handgun, Backpack Mistblower                  | 5% to 10% by volume  | partial coverage        |

<sup>1</sup>Where repeat applications are necessary, do not exceed 8.0 quarts per acre per year.

<sup>2</sup>For low volume directed spray applications, coverage should be uniform with at least 50 percent of the foliage contacted. For best results, coverage of the top one-half of the plant is important.

In forestry site preparation and utility rights-of-way applications, this product requires use with a surfactant such as a non-ionic surfactant containing greater than 80 percent active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Use higher rates of this product within the recommended rate ranges for control or partial control of woody brush, trees and hard-to-control perennial herbaceous weeds. For best results, apply to actively growing woody brush and trees after full leaf expansion and before fall color and leaf drop. Use increased rates within the recommended rate range to control of perennial herbaceous weeds from emergence up to the appearance of seedheads, flowers or berries appear. Use lower rates within the recommended rate range to control annual herbaceous weeds and actively growing perennial herbaceous weeds after seedheads, flowers or berries appear. Apply to foliage of actively growing annual herbaceous weeds anytime after emergence.

### Tank Mixtures

This product may be used in tank mix combination with other herbicide products to broaden the spectrum of vegetation controlled. When tank mixing, read and observe applicable use directions, precautions and limitations on the respective product labels. Use according to the most restrictive precautionary statements for each product on the mixture. Any recommended rate of this product may be used in a tank mix.

**Note:** For forestry site preparation, make sure the tank mix product is approved for use prior to planting the desired species. Observe planting interval restrictions. For side trimming treatments in utility rights-of-way, tank mixtures with Arsenal 2WSL herbicide are not recommended. For side trimming treatments, it is recommended that this product be used alone as recommended, or as a tank mix with Garlon.

| Product                                | Broadcast Rate          | Use Sites                                |
|----------------------------------------|-------------------------|------------------------------------------|
| Arsenal Applicators Concentrate        | 2 to 16 fl oz/acre      | Forestry site preparation                |
| Oust                                   | 1 to 4 oz/acre          | Forestry site preparation, utility sites |
| Garlon 3A <sup>1</sup>                 | 1 to 4 qt/acre          | Forestry site preparation, utility sites |
| Garlon 4                               | 1 to 4 qt/acre          | Forestry site preparation, utility sites |
| Arsenal 2WSL                           | 2 to 32 fl oz/acre      | Utility sites                            |
| <b>Spray-to-Wet Rates</b>              |                         |                                          |
| Arsenal Applicators Concentrate        | 1/32% to 1/2% by volume | Forestry site preparation                |
| Arsenal 2WSL                           | 1/32% to 1/2% by volume | Utility sites                            |
| <b>Low Volume Directed Spray Rates</b> |                         |                                          |
| Arsenal Applicators Concentrate        | 1/8% to 1/2% by volume  | Forestry site preparation                |
| Arsenal 2WSL                           | 1/8% to 1/2% by volume  | Utility sites                            |

<sup>1</sup>Ensure that Garlon 3A is thoroughly mixed with water before adding this product. Agitation is required while mixing this product with Garlon 3A to avoid compatibility problems.

For control of herbaceous weeds, use the lower recommended tank mixture rates. For control of dense stands or difficult-to-control woody brush and trees, use the higher recommended rates.

### Forestry Conifer and Hardwood Release

#### Directed Sprays and Selective Equipment

This product may be applied as a directed spray or by using selective equipment in forestry conifer and hardwood sites, including Christmas tree plantations and silvicultural nurseries. This product requires use with a surfactant. Use only surfactants that are approved for conifer release and specified on the surfactant label as safe for use in conifer release (pine release). Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

**Tank Mixing:** In hardwood plantations, tank mixtures with Oust may be used. In pine plantations, tank mixtures with Garlon 4 or Arsenal AC may be used. Comply with all site restrictions, forestry species limitations, and precautions on the tank mix product labels.

Avoid contact of spray drift, mist or drips with foliage, green bark or non-woody surface roots of desirable plant species. See "Application Equipment and Techniques" section of this label for specific recommendations and precautions.

**Spray-to-Wet Applications:** Use a 2 percent spray solution to control undesirable woody brush and trees. To control herbaceous weeds, use a 1 to 2 percent spray solution.

**Low Volume Directed Spray Applications:** Use a 5 to 10 percent spray solution. Coverage should be uniform with at least 50 percent of the foliage contacted. Coverage of the top one-half of the unwanted vegetation is important.

**Broadcast Applications:** For equipment calibrated for broadcast applications, use 1 1/2 to 7 1/2 quarts of this product per acre. Apply in 10 to 60 gallons of clean water per acre. Shielded application equipment may be used to avoid contact of the spray solution with desirable plants. Shields should be adjusted to prevent spray contact with the foliage of green bark of desirable vegetation.

**Wiper Application Equipment:** See the "Selective Equipment" section of this label for equipment and application rate recommendations.

### Broadcast Application

**Note:** Except where specifically recommended below, make broadcast applications of this product only where conifers have been established for more than one year.

**Broadcast application must be made after formation of final conifer resting buds in the fall or prior to initial bud swelling in the spring.**

Injury may occur to conifers treated for release, especially where spray patterns overlap or the higher rates are applied. Damage can be accentuated if applications are made when conifers are actively growing, or are under stress from drought, flood water, improper planting, insects, animal damage or diseases.

This product requires use with a surfactant. Use a surfactant that is labeled/recommended for use in over-the-top release applications. Use of this product without a surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

**For release of the following conifer species outside the Southeastern United States:**

Douglas fir (*Pseudotsuga menziesii*)  
Fir (*Abies* species)  
Hemlock <sup>†</sup> (*Tsuga* species)  
Pines <sup>†</sup> (*Pinus* species)  
Redwood, California <sup>†</sup> (*Sequoia* species)

<sup>†</sup> Includes all species except loblolly pine, longleaf pine, shortleaf pine or slash pine.

<sup>†</sup> Use of a surfactant is not recommended for release of hemlock species or California redwood. In mixed conifer stands, injury to these species may result if a surfactant is used.

**Application Rate for Conifer Release:** Apply 3/4 to 1 1/2 quarts per acre as a broadcast spray. In Maine and New Hampshire, up to 2 1/4 quarts per acre of this product may be used for the control and suppression of difficult-to-control hardwood species.

To release Douglas fir, and pine and spruce species at the end of the first growing season (except in California), apply 3/4 to 1 1/8 quarts per acre of this product. Make sure that all conifers are well hardened off.

**Note:** For release of Douglas fir with this product or recommended tank mixtures, a nonionic surfactant recommended for over-the-top foliar spray may be used. To avoid possible conifer injury, nonionic surfactants may be used at 2 fluid ounces per acre at elevations above 1500 feet, or 1 fluid ounce per acre in the coastal range or at elevations below 1500 feet. Use of surfactant rates exceeding those listed above may result in unacceptable conifer injury and are not recommended. Make sure that the nonionic surfactant has been adequately tested for safety to Douglas fir before use.

**Tank Mixtures with Oust:** To release jack pine, white pine and white spruce, apply 3/4 to 1 1/2 quarts of this product with 1 to 3 ounces (1 to 1 1/2 ounces for white pine) of Oust per acre. Make applications to actively growing weeds as a broadcast spray over the top of established conifers. Applications at these rates should be made after formation of conifer resting buds in the late summer or fall.

**Tank Mixtures with Arsenal Applicators Concentrate:** This product may be tank mixed with Arsenal Applicators Concentrate for release of Douglas fir. Tank mix 3/4 to 1 1/8 quarts of this product with 2 to 6 fluid ounces of Arsenal Applicators Concentrate per acre. For release of balsam fir and red spruce, apply a mixture of 1 1/2 quarts of this product with 1 to 2 1/2 fluid ounces of Arsenal Applicators Concentrate per acre.

In Maine and New Hampshire for the release of red pine, balsam fir, red spruce, white spruce, Norway spruce, and black spruce with dense tough-to-control brush and where maples make up a large component of the undesirable trees, up to 2 1/4 quarts per acre of this product may be tank mixed with 1 to 2 1/2 fluid ounces per acre of Arsenal Applicators Concentrate herbicide and applied as a broadcast spray.

**Tank mixtures with Arsenal Applicators Concentrate and Oust or Oust XP Herbicides:** In Maine and New Hampshire for release of red pine, balsam fir, red spruce, white spruce, Norway spruce and black spruce with heavy grass and herbaceous weed densities, tough-to-control brush and where maples make up a large component of the undesirable trees up to 2 1/4 quarts per acre of this product may be tank mixed with 1 to 2.5 fluid ounces per acre of Arsenal Applicators Concentrate and 1 to 3 oz of Oust or Oust XP herbicides and applied as a broadcast spray.

**For release of the following conifer species in the Southeastern United States:**

Loblolly pine (*Pinus taeda*)  
Eastern white pine (*Pinus strobus*)  
Shortleaf pine (*Pinus echinata*)  
Slash pine (*Pinus elliotii*)  
Virginia pine (*Pinus virginiana*)  
Longleaf pine (*Pinus palustris*)

Apply 1 1/8 to 1 7/8 quarts of this product per acre as a broadcast spray during late summer or early fall after the conifers have hardened off. For applications at the end of the first growing season, use 3/4 quart of this product alone or in a recommended tank mixture.

**Tank Mixtures with Arsenal Applicators Concentrate:** For conifer release, apply 3/4 to 1 1/2 quarts of this product with 2 to 16 fluid ounces of Arsenal Applicators Concentrate per acre as a broadcast spray. Use only on conifer species that are labeled for over-the-top spray for both products. Use the higher recommended rates for dense tough-to-control wood brush and trees.

Read and observe label claims, cautionary statements and all information on the labels of each product used in these tank mixtures. Use according to the most restrictive precautionary statements for each product in the mixture.

### Herbaceous Release

When applied as directed, this product plus listed residual herbicides provides postemergence control of the annual weeds and control or suppression of the perennial weeds listed in this label, and residual control of the weeds listed in the residual herbicide label. Make applications to actively growing weeds as a broadcast spray over the top of labeled conifers.

**Tank Mixtures with Oust:** To release loblolly pines, tank mix 12 to 18 fluid ounces of this product with 2 to 4 ounces of Oust per acre.

To release slash pines, tank mix 9 to 12 fluid ounces of this product with 2 to 4 ounces of Oust per acre.

In Maine and New Hampshire for release of red pine, balsam fir, red spruce, white spruce, Norway spruce, and black spruce with heavy grass and herbaceous weeds infesting the site, up to 2 1/4 quarts per acre of this product may be tank mixed with 1 to 3 oz of Oust herbicide or Oust XP herbicide to control grass, herbaceous weeds and woody brush, and applied as a broadcast spray.

For tank mixtures with Oust use a surfactant that is labeled/ recommended for use in over-the-top herbaceous release applications. Use of this product without a surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Weed control may be reduced if water volumes exceed 25 gallons per acre for these treatments.

**Tank Mixture with Atrazine:** To release Douglas fir, apply 3/4 quart of this product with 4 pounds a.i. of atrazine per acre. Apply only over Douglas fir that has been established for at least one full growing season. Apply in the early spring, usually mid-March through early April. Injury will occur if applications are made after bud swell in the spring. For this use, do not add surfactant to the tank mixture.

Always read and follow the manufacturer's label for all herbicides and surfactants used.

### Wetland Sites

This product may be used in and around water (aquatic areas) and wetlands found in forestry and in power, telephone and pipeline rights-of-way sites, including where these sites are adjacent to or surrounding domestic water supply reservoirs, supply streams, lakes and ponds. Read and observe the following before making applications in and around water.

Consult local public water control authorities before applying this product in and around public water. Permits may be required to treat in such areas.

There is no restriction on the use of treated water for irrigation, recreation or domestic purposes.

**Note:** Do not apply this product directly to water within 1/2 mile up-stream of an active potable water intake in flowing water (i.e., river, stream, etc.) or within 1/2 mile of an active potable water intake in a standing body of water such as a lake, pond or reservoir. To make aquatic applications around and within 1/2 mile of active potable water intakes, the water intake must be turned off for a minimum period of 48 hours after application. These aquatic applications may be made ONLY in those cases where there are alternative water sources or holding ponds which would permit the turning off of an active potable water intake for a minimum period of 48 hours after the application. This restriction does not apply to intermittent inadvertent overspray of water in terrestrial use sites.

Do not spray open bodies of water where woody brush, trees and herbaceous weeds do not exist. The maximum application rate of 3 3/4 quarts per acre must not be exceeded in a single over-water broadcast application except as follows, where any recommended rate may be applied:

- Stream crossings in utility right-of-way.
- Where applications will result in less than 20 percent of the total water area being treated.

### Wildlife Habitat Restoration and Management Areas

This product is recommended for the restoration and/or maintenance of native habitat and in wildlife management areas.

**Habitat Restoration and Maintenance:** When applied as directed, exotic and other undesirable vegetation may be controlled in habitat management areas. Applications may be made to allow recovery of native plant species, to open up water to attract waterfowl, and for similar broad-spectrum vegetation control requirements in habitat management areas. Spot treatments may be made to selectively remove unwanted plants for habitat enhancement. For spot treatments, care should be exercised to keep spray off of desirable plants.

**Wildlife Food Plots:** This product may be used as a site preparation treatment prior to planting wildlife food plots. Apply as directed to control vegetation in the plot area. Any wildlife food species may be planted after applying this product, or native species may be allowed to reinfest the area. If tillage is needed to prepare a seedbed, wait 7 days after applying this product before tilling to allow for maximum effectiveness.

## Wiper Applications

For wick or wiper applications, mix 1 gallon of this product with 2 gallons of clean water to make a 33 percent solution. Addition of a nonionic surfactant at a rate of 10 percent by volume of total herbicide solution is recommended.

Wiper applications can be used to control or suppress annual and perennial weeds listed on this label. In heavy weed stands, a double application in opposite directions may improve results. See the "Weed Controlled" section in this label for recommended timing, growth stage and other instructions for achieving optimum results

## Cut Stump Application

Woody vegetation may be controlled by treating freshly cut stumps of trees and resprouts with this product. Apply this product using suitable equipment to ensure coverage of the entire cambium. Cut vegetation close to the soil surface. **Apply a 50 to 100 percent solution of this product to freshly cut surface immediately after cutting.** Delay in applying this product may result in reduced performance. For best results, trees should be cut during periods of active growth and full leaf expansion.

When used according to directions for cut stump application, this product will **control, partially control or suppress** most woody brush and tree species, some of which are listed below:

| Common Name    | Scientific Name                |
|----------------|--------------------------------|
| Alder          | <i>Alnus spp.</i>              |
| Coyote brush † | <i>Baccharis consanguinea</i>  |
| Dogwood †      | <i>Cornus spp.</i>             |
| Eucalyptus     | <i>Eucalyptus spp.</i>         |
| Hickory †      | <i>Carya spp.</i>              |
| Madrone        | <i>Arbutus menziesii</i>       |
| Maple †        | <i>Acer spp.</i>               |
| Oak            | <i>Quercus spp.</i>            |
| Poplar †       | <i>Populus spp.</i>            |
| Reed, giant    | <i>Arundo donax</i>            |
| Salt cedar     | <i>Tamarix spp.</i>            |
| Sweet gum †    | <i>Liquidambar styraciflua</i> |
| Sycamore †     | <i>Platanus occidentalis</i>   |
| Tan oak        | <i>Lithocarpus densiflorus</i> |
| Willow         | <i>Salix spp.</i>              |

† This product is not approved for this use on these species in the state of California.

## Injection and Frill Applications

Woody vegetation may be controlled by injection or frill application of this product. Apply this product using suitable equipment which must penetrate into living tissue. Apply the equivalent of 1 ml of this product per 2 to 3 inches of trunk diameter. This is best achieved by applying 25 to 100 percent concentration of this product either to a continuous frill around the tree or as cuts evenly spaced around the tree below all branches. As tree diameter increases in size, better results are achieved by applying dilute material to a continuous frill or more closely spaced cuttings. Avoid application techniques that allow runoff to occur from frill or cut areas in species that exude sap freely after frills or cutting. In species such as these, make frill or cut at an oblique angle so as to produce a cupping effect and use undiluted material. For best results, applications should be made during periods of active growth and full leaf expansion.

This treatment will control the following woody species:

| Common Name | Scientific Name                |
|-------------|--------------------------------|
| Oak         | <i>Quercus spp.</i>            |
| Poplar      | <i>Populus spp.</i>            |
| Sweet gum   | <i>Liquidambar styraciflua</i> |
| Sycamore    | <i>Platanus occidentalis</i>   |

This treatment will suppress the following woody species:

| Common Name | Scientific Name        |
|-------------|------------------------|
| Black gum † | <i>Nyssa sylvatica</i> |
| Dogwood     | <i>Cornus spp.</i>     |
| Hickory     | <i>Carya spp.</i>      |
| Maple, red  | <i>Acer rubrum</i>     |

† This product is not approved for this use on this species in the state of California.

## Release of Bermudagrass or Bahiagrass on Noncrop Sites

### Release Of Dormant Bermudagrass And Bahiagrass

When applied as directed, this product will provide control or suppression of many winter annual weeds and tall fescue for effective release of dormant bermudagrass or bahiagrass. Make applications to dormant bermudagrass or bahiagrass.

For best results on winter annuals, treat when weeds are in an early growth stage (below 6 inches in height) after most have germinated. For best results on tall fescue, treat when fescue is in or beyond the 4 to 6-leaf stage.

## Weeds Controlled

Rate recommendations for control or suppression of winter annuals and tall fescue are listed below.

Apply the recommended rates of this product in 10 to 25 gallons of water per acre plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient.. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

### Weeds Controlled or Suppressed <sup>†</sup>

**Note:** C = Controlled; S = Suppressed

| Weed Species                                       | Rate<br>(Fluid Ounces Per Acre) |   |    |    |    |    |
|----------------------------------------------------|---------------------------------|---|----|----|----|----|
|                                                    | 6                               | 9 | 12 | 18 | 24 | 48 |
| Barley, little<br><i>Hordeum pusillum</i>          | S                               | C | C  | C  | C  | C  |
| Bedstraw, catchweed<br><i>Galium aparine</i>       | S                               | C | C  | C  | C  | C  |
| Bluegrass, annual<br><i>Poa annua</i>              | S                               | C | C  | C  | C  | C  |
| Chervil<br><i>Chaerophyllum tainturieri</i>        | S                               | C | C  | C  | C  | C  |
| Chickweed, common<br><i>Stellaria media</i>        | S                               | C | C  | C  | C  |    |
| Clover, crimson<br><i>Trifolium incarnatum</i>     | •                               | S | S  | C  | C  | C  |
| Clover, large hop<br><i>Trifolium campestre</i>    | •                               | S | S  | C  | C  | C  |
| Speedwell, corn<br><i>Veronica arvensis</i>        | S                               | C | C  | C  | C  | C  |
| Fescue, tall<br><i>Festuca arundinacea</i>         | •                               | • | •  | •  | S  | S  |
| Geranium, Carolina<br><i>Geranium carolinianum</i> | •                               | • | S  | S  | C  | C  |
| Henbit<br><i>Lamium amplexicaule</i>               | •                               | S | C  | C  | C  | C  |
| Ryegrass, Italian<br><i>Lolium multiflorum</i>     | •                               | • | S  | C  | C  | C  |
| Vetch, common<br><i>Vicia sativa</i>               | •                               | • | S  | C  | C  | C  |

<sup>†</sup>These rates apply only to sites where an established competitive turf is present.

## Release Of Actively Growing Bermudagrass

**NOTE:** Use only on sites where bahiagrass or bermudagrass are desired for ground cover and some temporary injury or yellowing of the grasses can be tolerated.

When applied as directed, this product will aid in the release of bermudagrass by providing control of annual species listed in the "Weeds Controlled" section in this label, and suppression or partial control of certain perennial weeds.

For control or suppression of those annual species listed in this label, use 3/4 to 2 1/4 pints of this product as a broadcast spray in 10 to 25 gallons of spray solution per acre, plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient.. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information. Use the lower rate when treating annual weeds below 6 inches in height (or length of runner in annual vines). Use the higher rate as size of plants increases or as they approach flower or seedhead formation.

Use the higher rate for partial control or longer-term suppression of the following perennial species. Use lower rates for shorter-term suppression of growth.

|               |                              |
|---------------|------------------------------|
| Bahiagrass    | Johnsongrass <sup>1</sup>    |
| Dallisgrass   | Trumpet creeper <sup>2</sup> |
| Fescue (tall) | Vaseygrass                   |

<sup>1</sup> Johnsongrass is controlled at the higher rate.

<sup>2</sup> Suppression at the higher rate only.

Use only on well-established bermudagrass. Bermudagrass injury may result from the treatment but regrowth will occur under moist conditions. Repeat applications in the same season are not recommended, since severe injury may result.

### **Bahiagrass Seedhead and Vegetative Suppression**

When applied as directed in the "Noncrop Sites" section in this label, this product will provide significant inhibition of seedhead emergence and will suppress vegetative growth for a period of approximately 45 days with single applications and approximately 120 days with sequential applications.

Apply this product 1 to 2 weeks after full green-up of bahiagrass or after the bahiagrass has been mowed to a uniform height of 3 to 4 inches. Applications must be made prior to seedhead emergence. Apply 5 fluid ounces per acre of this product in 10 to 25 gallons of water per acre, plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information.

Sequential applications of this product plus nonionic surfactant may be made at approximately 45-day intervals to extend the period of seedhead and vegetative growth suppression. For continued vegetative growth suppression, sequential applications must be made prior to seedhead emergence.

Apply no more than 2 sequential applications per year. As a first sequential application, apply 3 fluid ounces of this product per acre plus nonionic surfactant. A second sequential application of 2 to 3 fluid ounces per acre plus nonionic surfactant may be made approximately 45 days after the last application.

### **Annual Grass Growth Suppression**

For growth suppression of some annual grasses, such as annual ryegrass, wild barley and wild oats growing in coarse turf on roadsides or other industrial areas, apply 3 to 4 ounces of this product in 10 to 40 gallons of water per acre plus a surfactant such as a non-ionic surfactant containing 80% or greater active ingredient. Use of this product without surfactant will result in reduced herbicide performance. Refer to the "Mixing and Application Instructions" section of this label and the surfactant manufacturer label for more information. Applications should be made when annual grasses are actively growing and before the seedheads are in the boot stage of development. Treatments made after seedhead emergence may cause injury to the desired grasses.

### **Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

### **Warranty Disclaimer**

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

### **Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. To the fullest extent permitted by law, all such risks shall be assumed by buyer.

### **Limitation of Remedies**

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- (1) Refund of purchase price paid by buyer or user for product bought, or
- (2) Replacement of amount of product used.

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. To the fullest extent permitted by law, in no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

<sup>®</sup>Trademark of Dow AgroSciences LLC

**Produced for**  
**Dow AgroSciences LLC**  
9330 Zionsville Road  
Indianapolis, IN 46268

Label Code: D02-148-005  
Replaces Label: D02-148-004  
LOES Number: 010-01471

EPA-accepted 07/13/06

### **Revisions:**

1. Updated Storage and Disposal