

432-1516

1/17/2014

1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Richard M. Gorrell  
Bayer Environmental Science  
2 T.W. Alexander Drive  
Research Triangle Park, NC 27709

JAN 17 2014

Subject: Notification; Per PR-Notice 98-10  
Esplanade 200 SC Herbicide  
EPA Reg. No. 432-1516  
Date Submitted: January 14, 2014

Dear Mr. Gorrell:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated January 14, 2014 for the product referenced above. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact me at (703) 306-0415 or [davis.kable@epa.gov](mailto:davis.kable@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Kable Bo Davis", written over a horizontal line.

Kable Bo Davis  
Product Manager 25  
Herbicide Branch  
Registration Division (7505P)

2/12



United States  
Environmental Protection Agency  
Washington, DC 20460

 Registration  
 Amendment  
 Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 432-1516	2. EPA Product Manager Kable Bo Davis	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Esplanade 200 SC Herbicide	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) Bayer Environmental Science 2 T.W. Alexander Drive RTP, NC 27709 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated <b>NOTIFICATION</b>
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

**JAN 17 2014**

**Explanation:** Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance of PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and it may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

-Minor change to the label as highlighted

### Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt.      No. per container	If "Yes" Package wgt      No. per container

3. Location of Net Contents Information  
 Label  Container

4. Size(s) Retail Container  
1x55 gallon drum

5. Location of Label Directions

6. Manner in Which Label is Affixed to Product  
 Lithograph Paper glued Stenciled  Other \_\_\_\_\_

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Richard M. Gorrell	Title Manager, Registrations	Telephone No. (Include Area Code) 919-549-2423
----------------------------	---------------------------------	---

<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature <i>Richard M. Gorrell</i>	3. Title Manager, Registrations	
4. Typed Name Richard M. Gorrell	5. Date January 14, 2014	



4/12

# NOTIFICATION

JAN 17 2014

GROUP 29 HERBICIDE

# Esplanade 200 SC

Preemergent Herbicide for the Control of Annual Grasses and Broadleaf Weeds in Non-Residential Non-Crop Areas, Railroad and Rail Yards, Managed Roadsides, Fence Rows, Utilities, Hardscapes, Industrial, Municipal, and Government Sites

**ACTIVE INGREDIENT:**

Indaziflam (CAS No: 730979-19-8).....19.05%

**OTHER INGREDIENTS:**.....80.95%

**TOTAL**.....100.00%

**EPA Reg. No. 432-1516**

**EPA Est.**

Contains 1.67 pounds of indaziflam per gallon

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

For **MEDICAL** and **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577  
For **PRODUCT USE** Information Call 1-800-331-2867

### FIRST AID


<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If on skin:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p><b>For MEDICAL Emergencies Call 24 Hours A Day 1-800-334-7577</b></p> <p>Have the product container or label with you when calling a poison control center or doctor or going for treatment.</p>	

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed, absorbed through the skin or inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. All mixers, loaders, applicators and other handlers must wear long-sleeved shirt, long pants, shoes plus socks, and water proof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using toilet. Remove and wash contaminated clothing before reuse.



## ENVIRONMENTAL HAZARDS

This Product is toxic to fish, aquatic invertebrates, and plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean watermark. Do not contaminate water when disposing of rinsate or washwater. This product may impact water through spray drift or runoff. Follow directions for use to avoid spray drift and runoff. A level well maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential of this product entering water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

**Ground Water Advisory:** This pesticide has properties and characteristics associated with chemicals detected in groundwater. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

**Surface Water Advisory:** This pesticide may impact water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application.

## DIRECTIONS FOR USE

**It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.  
Read the entire label before using this product**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**SHAKE CONTAINER WELL BEFORE USING.**

## PRODUCT INFORMATION

Esplanade 200 SC is a selective, preemergent, alkyazine herbicide for control of many annual grasses and broadleaf weeds in railroad, roadside, hardscapes, industrial areas, utilities, airports, government and military installations, managed areas (petroleum tank farms, pumping stations, storage areas, rail and utility rights-of-way, utility substations, lumberyards, around farm buildings, non-irrigation ditch banks, fence rows, manufacturing sites, office buildings, educational facilities, parking lots, and under asphalt or concrete as part of site preparation).

Esplanade 200 SC controls weeds by reducing the emergence of seedlings through inhibition of cellulose biosynthesis (CB Inhibitor). Necrosis or yellowing may also be observed if the herbicide is applied to herbaceous tissue such as leaves and green stems of susceptible plants. The herbicide needs to be activated prior to weed germination for most effective control. For maximum activity against germinating weeds, Esplanade 200 SC requires rainfall (minimum 0.25 inches) within several weeks after application to activate the herbicide.

Esplanade 200 SC has minimal post emergent activity and generally does not control weeds that have emerged. A post emergent herbicide such as Finale® Herbicide may be mixed with Esplanade 200 SC to control existing weeds. Esplanade 200 SC does not control tubers, rhizomes, and woody vegetation.

Esplanade 200 SC can be applied to terrestrial non-crop sites and unimproved turf sites that contain areas of casual water of a temporary nature as a result of surface water collecting in equipment wheel ruts or in other depressions created by management activities.

Esplanade 200 SC may only be applied by ground equipment only.

## USE RESTRICTIONS

- Do not apply directly to water or to soil where standing water is present except as specified on this label.
- Do not apply in or on irrigation ditches/canals including the outer banks.
- Do not contaminate water intended for irrigation and domestic use.
- Do not treat or allow spray drift or runoff to fall into irrigation ditches/canals or other channels that carry water; that may be used for irrigation purposes.
- Do not exceed 7 fl oz per acre of Esplanade 200 SC in a single application for all Industrial Vegetation Management.
- Do not exceed 10 fl oz per acre of Esplanade 200 SC for all Industrial Vegetation Management applications within a calendar year or in a 12-month period from the previous application.
- Do not apply Esplanade 200 SC to newly seeded turf.
- Do not apply Esplanade 200 SC through an irrigation or chemigation system.
- Do not apply by air.

- Do not apply or otherwise permit this product or sprays containing this product to come into contact with any non-target crop or desirable plants.
- Do not make applications when circumstances favor movement from treatment sites.
- Do not apply to frozen or snow covered ground.
- Do not graze or feed forage, hay, or straw from treated areas to livestock.
- Do not use on residential lawns, golf courses, sod farms, or production and landscape ornamentals.
- Esplanade 200 SC is not for sale, distribution, or use in Nassau County or Suffolk County in New York State.

**USE PRECAUTIONS**

- Applications made to areas where runoff water flows onto agricultural land may injure crops.
- Applications made during periods of intense rainfall, to soils saturated with water, or soils through which rainfall will not readily penetrate may result in runoff and movement of Esplanade 200 SC.
- Treated soil should be left undisturbed to reduce the potential for Esplanade 200 SC movement by soil erosion, by wind, or water.
- Applications should be made only when there is little or no risk of spray drift or movement of applied product into sensitive areas. Sensitive areas are defined as bodies of water (ponds, lakes, rivers, and streams), habitats of endangered species and non-labeled agricultural crop areas. Refer to the Spray Drift Management section of this label for more details.

**APPLICATION INFORMATION**

Apply Esplanade 200 SC with a properly calibrated sprayer according to the manufacturer's directions and check periodically to be certain that the equipment is working properly prior to each use. Uniform application is essential for satisfactory weed control. Avoid overlap. Shut off spray booms while starting, turning, slowing, or stopping to avoid off-target application.

When spraying close or next to ponds, lakes, rivers, and streams be cognizant of keeping the spray solution from reaching the water.

For all applications, follow these guidelines: use spray volumes of 10-100 gallons per acre, spray boom height and spray pressures as low as practical, use coarse droplet producing nozzle tips, use drift control additives and shielded sprayers where practical, and spray when wind speed is low. See the Spray Drift Management section for more details. The use of a hand-held or backpack sprayer is allowed, especially when treating smaller areas. The water volume and use rates are the same on a given area as if treating with a much larger boom sprayer. When using a hand-held or backpack sprayer, do not exceed the use rate restrictions stated on this label.

**MIXING INSTRUCTIONS**

Ensure that the application equipment has been thoroughly cleaned from previous use before using to apply Esplanade 200 SC. Fill the spray tank with 1/2 of the required volume of water prior to the addition of Esplanade 200 SC. Add the proper amount of Esplanade 200 SC, and then add the rest of the water. Maintain sufficient agitation to ensure an adequate spray mixture during application. If Esplanade 200 SC is to be applied in a tank mixture with other pesticides, add the appropriate amounts of the tank mix partners in the following order: (a) products in water-soluble packaging (WSP), (b) WP, (c) WG or other dry flowables, (d) fertilizers, (e) Esplanade 200 SC, (f) other aqueous suspension products (SC), (g) liquid flowables, (h) emulsifiable concentrates and other organic-solvent based formulations. Continue to fill the tank with water to the desired volume while agitating. **Maintain sufficient agitation during application to ensure a uniform spray mixture.**

**Resuspending Esplanade 200 SC in Spray Solution:** Like other suspension concentrates (SC), Esplanade 200 SC will settle if left standing without agitation. Re-agitate the spray solution before application.

**COMPATIBILITY TESTING WITH OTHER PESTICIDES**

A compatibility test must be conducted with any potential tank mix partner with Esplanade 200 SC. Using a clear container, conduct the test as described below:

Fill the container three-quarters full with water.

1. Add the appropriate amount of tank mix partner in the following order: (a) WP (b) dry flowable (c) Esplanade 200 SC (d) aqueous suspensions, (e) flowables, (f) liquids and (g) solutions and emulsifiable or liquid concentrates. Shake or gently stir after each addition to mix thoroughly.
2. After adding all ingredients, let the mixture stand for 15 minutes and look for separation, large flakes, precipitates, gels, and heavy oily film or other signs of incompatibility.
3. If the compatibility test shows signs of incompatibility, do not tank mix the product tested with Esplanade 200 SC.

**Spray Drift Management**

Spray equipment and weather affect spray drift. Consider all factors when making application decisions. Where states have more stringent regulations, they must be observed. Avoiding spray drift is the responsibility of the applicator. To reduce the potential for drift, the ground application equipment must be set to apply COARSE or greater droplets (i.e., ASABE Standard 572.1) with corresponding spray pressure. Use high flow rate nozzles to apply the highest practical spray volume. With most nozzle types, narrower spray angles

produce larger droplets. Follow the nozzle manufacturer's directions on pressure, orientation, spray volume, etc., in order to minimize drift and optimize coverage and control.

**Sensitive Areas**

Sensitive areas are defined as bodies of water (ponds, lakes, rivers, and streams), wetlands, habitats of endangered species and non-labeled agricultural crop areas. Applicators must take all precautions necessary to keep spray drift from reaching sensitive areas.

Only apply this product when the potential for drift to adjacent sensitive areas is minimal (e.g. when wind is blowing away from the sensitive areas). The applicator is responsible for considering all these factors when making decisions.

**Wind**

Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing. Many factors influence spray drift potential including droplet size, equipment type, and local terrain. Drift potential increases if wind is in excess of 10 mph, gusty, or below 2 mph (due to inversion potential). Always make applications when there is some air movement to determine the direction and distance of possible spray drift. The applicator should be familiar with local conditions and how it may influence spray drift.

**Temperature Inversion**

A surface temperature inversion (i.e., increasing temperature with increasing altitude) greatly increases the potential for drift. Avoid application when conditions are favorable to inversion. Presence of ground fog is a good indicator of a surface temperature inversion.

**Controlling Droplet Size**

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that still provide sufficient coverage and control. Uniform spray coverage is important to maximize weed control. Applying larger droplets will reduce drift potential, but will not prevent drift if applications are made improperly or under unfavorable environmental conditions such as wind speed, temperature and humidity, and temperature inversion situations.

Spray volume, pressure, and nozzle selection are all important for reducing drift. Select a high flow rate nozzle to apply the highest practical spray volume. High flow rate nozzles produce larger droplets. Use lower spray pressures within the recommended range for the nozzle. If a higher flow rate is needed, increase the nozzle size instead of increasing pressure. Lower spray pressures produce larger droplets. Also, consider using low-drift nozzles.


Set the boom and make applications at the lowest height that safely permits uniform coverage of the soil and minimizes droplet evaporation. Avoid application if wind conditions are gusty. Local terrain may influence wind patterns. The applicator should be familiar with local conditions and understand how they may impact spray drift.

**Drift Control Additive**

Drift control additive may also be used with most spray equipment to reduce the potential for drift. When using a drift control additive, read and follow all directions on the additive label.

**Shielded Sprayers**

Shielding the boom or individual nozzles may also reduce the potential for drift. However, it is the responsibility of the applicator to verify that the shield does not interfere with uniform spray coverage.







Grasses and Sedges Controlled	
Annual bluegrass	<i>Poa annua</i>
Annual brome	<i>Bromus</i> spp.
Barnyardgrass, common	<i>Echinochloa crus-galli</i>
Cheatgrass	<i>Bromus secalinus</i>
Crabgrass	<i>Digitaria</i> species
Crabgrass, Henry	<i>Digitaria adscendens</i>
Crabgrass, Large/Hairy	<i>Digitaria sanguinalis</i>
Crabgrass, Smooth	<i>Digitaria ischaemum</i>
False chamomile	<i>Matricaria maritime</i>
Foxtail brome	<i>Bromus rubens</i>
Foxtail, Giant	<i>Setaria faberi</i>
Foxtail, Green	<i>Setaria viridis</i>
Foxtail, Yellow	<i>Pennisetum glaucum</i>
Goosegrass	<i>Eleusine indica</i>
Guineagrass	<i>Panicum maximum</i>
Medusahead	<i>Taeniatherum caput-medusae</i>
Mouse barley	<i>Hordeum murinum</i>
Rice flatsedge	<i>Cyperus iria</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>
Ryegrass, Perennial	<i>Lolium perenne</i>
Sedge, annual	<i>Cyperus</i> spp.
Sprangletop	<i>Leptochloa</i> spp.
Tufted lovegrass	<i>Eragrostis pectinacea</i>
Weeds Suppressed	
Black medic	<i>Medicago lupulina</i>
Black mustard	<i>Brassica nigra</i>
False chamomile	<i>Matricaria maritime</i>
London rocket	<i>Sisymbrium irio</i>
Nutsedge, purple	<i>Cyperus rotunda</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Prickly lettuce	<i>Lactuca serriola</i>
Sesbania, hemp	<i>Sesbania exaltata</i>
Sida, prickly/teaweed	<i>Sida spinosa</i>
Southern brassbuttons	<i>Cotula australis</i>
Sunflower, common	<i>Helianthus</i> spp.
Vetch, purple	<i>Vicia benghalensis</i>
Wild carrot	<i>Daucus carota</i>
Woodsorrell, yellow	<i>Oxalis stricta</i>
Woodsorrel/Oxalis	<i>Oxalis</i> species

Use Sites for Esplanade 200 SC	Rate Range (fl oz/A)	Maximum Single Use Rate ( fl oz/A)	Maximum Total Yearly Rate (fl oz/A)
Rail and Rail Yards	3.5*-7	7	10
Managed Roadsides	3.5-7	7	10
Warm Season Turf Release	3.5-5	5	10
All other use sites listed	3.5-7	7	10

\*In Rail and Rail yard use sites, the 3.5 oz rate of Esplanade 200 SC should only be applied under low weed pressure in combination with another approved herbicide. This rate is not intended for stand-alone treatments.

**Bareground Applications for Non-Residential Non-Crop Sites**

Bareground is desired at many non-crop sites for reducing fire hazards, maintaining appropriate lines-of-site, and aesthetic considerations. Examples of sites include but are not limited to guardrails and some median strips near highways, hardscapes, parks, airports, utilities, government and military installations, around farm buildings, manufacturing sites, office buildings, educational facilities, parking lots, and managed areas. Esplanade 200 SC may be used alone for residual weed control or in tank mixture. Tank mixtures with post emergent herbicides help to control existing weeds. Observe use restrictions for all herbicides if a tank mixture is applied. Use-rates for bareground applications depend on the duration of weed control desired and the weed species listed on this label. Apply Esplanade 200 SC at 5-7 fl oz per acre. A repeat application can be made but not to exceed a total amount of 10 fl oz per acre per year.

**Restriction:** Applications to hardscapes (e.g. patios, paved parking lots, and walkways) may be made by spot application only.

**Railroads and Rail Yards**

Esplanade 200 SC may be used for preemergent residual control of certain weeds near railroad tracks, ballasts, and rail yards. Follow application instructions under **Bareground Applications** where bareground is the desired result. In situations where warm season turfgrass coverage is desired, such as at railroad crossings, follow use directions under the **Warm Season Turf Release** section of this label. Apply Esplanade 200 SC at 5-7 fl oz per acre. A repeat application can be made but not to exceed a total amount of 10 fl oz per acre per year.

**Warm Season Turf Release**

Esplanade 200 SC may be used to promote the growth of warm season grasses in areas where low maintenance vegetation or erosion control is desired. Established bermudagrass (*Cynodon dactylon*), centipedegrass (*Eremochloa ophiuroides*), bahiagrass (*Paspalum notatum*), buffalograss (*Buchloe dactyloides*), and Zoysiagrass (*Zoysia spp.*) are tolerant to Esplanade 200 SC at rates up to 5 fl oz per acre. Application of Esplanade 200 SC in the spring or fall to these grasses will control labeled weeds and allow low maintenance turf to develop. A repeat application can be made but not to exceed a total amount of 10 fl oz per acre per year. Cool season grasses such as Kentucky bluegrass (*Poa pratensis*), perennial ryegrass (*Lolium perenne*), and fescues (*Festuca sp*) are not tolerant to Esplanade 200 SC. Use Esplanade 200 SC on these grasses only when removal of these grasses is desired.

Esplanade 200 SC can inhibit the emergence of seed and damage newly emerged seedlings. Seeding into turf treated with Esplanade 200 SC should be delayed until at least **8 months** after application. Applications to newly seeded turf made sooner than **8 months** after emergence may significantly reduce stand establishment and turf vigor.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Protect ESPLANADE™ 200 SC from freezing temperatures.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous.

Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or reconditioning, if appropriate. Then puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**Greater than 5 gallons:**

Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling, if available or reconditioning, if appropriate or dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

